



TITLE:

THE ABELIANIZATION OF THE CONGRUENCE IA-AUTOMORPHISM GROUP OF A FREE GROUP(Methods of Transformation Group Theory)

AUTHOR(S):

Satoh, Takao

CITATION:

Satoh, Takao. THE ABELIANIZATION OF THE CONGRUENCE IA-AUTOMORPHISM GROUP OF A FREE GROUP(Methods of Transformation Group Theory). 数理解析研究所講究録 2006, 1517: 60-68

ISSUE DATE:

2006-10

URL:

<http://hdl.handle.net/2433/58715>

RIGHT:

THE ABELIANIZATION OF THE CONGRUENCE IA-AUTOMORPHISM GROUP OF A FREE GROUP

佐藤隆夫 (Takao Satoh)

東京大学大学院数理科学研究科 (The University of Tokyo)

Dedicated to Professor Kojun Abe on the occasion of his 60th birthday

ABSTRACT. In this paper we consider the abelianizations of some normal subgroups of the automorphism group of a finitely generated free group. Let F_n be a free group of rank n . For $d \geq 2$, we consider a group consisting the automorphisms of F_n which act trivially on the first homology group of F_n with $\mathbf{Z}/d\mathbf{Z}$ -coefficients. We call it the congruence IA-automorphism group of level d and denote it by $IA_{n,d}$. Let $IO_{n,d}$ be the quotient group of the congruence IA-automorphism group of level d by the inner automorphism group of a free group. In this paper we determine the abelianization of $IA_{n,d}$ and $IO_{n,d}$ for $n \geq 2$ and $d \geq 2$. Furthermore, for $n = 2$ and odd prime p , we compute the integral homology groups of $IA_{2,p}$ for any dimension.

1. INTRODUCTION

Let F_n be a free group of rank n , and $\text{Aut } F_n$ the automorphism group of the free group F_n . We denote the abelianization of F_n by H . The abelianization homomorphism $F_n \rightarrow H$ induces a surjective homomorphism $\rho : \text{Aut } F_n \rightarrow GL(n, \mathbf{Z})$. In this paper we consider the abelianization of the preimages of the congruence subgroups of $GL(n, \mathbf{Z})$ by ρ . For $n \geq 2$ and $d \geq 2$, let $GL(n, d)$ be the general linear group over $\mathbf{Z}/d\mathbf{Z}$, and $\pi_d : GL(n, \mathbf{Z}) \rightarrow GL(n, d)$ the natural homomorphism induced by the mod reduction d . We call the kernel $\Gamma(n, d)$ of π_d the congruence subgroup of $GL(n, d)$ of level d . Classically, the congruence subgroups $\Gamma(n, d)$ have been studied by many authors, and there is a broad range remarkable results of them. In particular, it is well known that for $n \geq 3$ and odd prime integer p , Lee and Szczarba [10, Theorem 1.1] determined the structure of the abelianization of the congruence subgroup $\Gamma(n, p)$. More precisely, they showed that it is isomorphic to the Lie algebra $\mathfrak{sl}_n(\mathbf{F}_p)$ of trace-zero matrices over \mathbf{F}_p as an $SL(n, \mathbf{F}_p)$ -module where \mathbf{F}_p is the finite field of order p .

2000 *Mathematics Subject Classification.* 20F28, 20F12, 20F14, 20F40, 16W25(Primary), 20F38, 57M05(Secondly).

Key words and phrases. the automorphism group of a free group, the Johnson homomorphism, Morita's trace.

In this paper we are also interested in the kernel IA_n of the natural map ρ . We call it the IA-automorphism group of F_n . Then we have an exact sequence

$$1 \rightarrow IA_n \rightarrow \text{Aut } F_n \xrightarrow{\rho} GL(n, \mathbf{Z}) \rightarrow 1.$$

This exact sequence plays important roles in the study of $\text{Aut } F_n$. Although Magnus [11] obtained a finitely many generating set of IA_n for $n \geq 3$, (See Subsection 2.1.) it is not known whether IA_n is finitely presented or not for $n \geq 4$. We remark that Krstić and McCool [9] showed that IA_3 is not finitely presentable.

For a group G , we denote the abelianization of G by G^{ab} . Cohen-Pakianathan [2, 3], Farb [4] and Kawazumi [8] independently showed that the abelianization IA_n^{ab} of IA_n is a free abelian group of rank $n^2(n-1)/2$. More precisely, it is isomorphic to $H^* \otimes_{\mathbf{Z}} \Lambda^2 H$ as a $GL(n, \mathbf{Z})$ -module where H^* is the dual group $\text{Hom}_{\mathbf{Z}}(H, \mathbf{Z})$ of H . We remark that the $GL(n, \mathbf{Z})$ -module structure of IA_n^{ab} is determined by using the first Johnson homomorphism of $\text{Aut } F_n$.

Here we consider subgroups of $\text{Aut } F_n$ which corresponds to the congruence subgroup of $GL(n, \mathbf{Z})$. Let $IA_{n,d}$ be the kernel of $\pi_d \circ \rho : \text{Aut } F_n \rightarrow GL(n, d)$, and call it the congruence IA-automorphism group of a free group F_n of level d . The first aim of this paper is to determine the structure of the abelianization of $IA_{n,d}$. Then our first result is

Theorem 1.1. *For $n \geq 2$ and $d \geq 2$,*

$$IA_{n,d}^{\text{ab}} \simeq (IA_n^{\text{ab}} \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z}) \bigoplus \Gamma(n, d)^{\text{ab}}.$$

In Section 3, we prove this theorem using the “extended” Johnson homomorphism, introduced by Kawazumi [8], constructed from the $\mathbf{Z}/d\mathbf{Z}$ -valued Magnus expansion of $\text{Aut } F_n$. Considering the result of Lee and Szczarba [10] stated above, we see that for any odd prime p , the abelianization of IA_n is isomorphic to $(\mathbf{Z}/p\mathbf{Z})^{\oplus \frac{1}{2}(n-1)(n^2+2n+2)}$ as an abelian group.

Next we consider the outer automorphism group of a free group and the images of IA_n and $IA_{n,d}$ by a natural projection. An automorphism ι of F_n is called an inner automorphism of F_n if there exists some element $y \in F_n$ such that $x' = y^{-1}xy$ for any $x \in F_n$. Then the group $\text{Inn } F_n$ of inner automorphisms of F_n is a normal subgroup of $\text{Aut } F_n$. Let $\text{Out } F_n$ be the quotient group $\text{Aut } F_n / \text{Inn } F_n$. The groups $\text{Inn } F_n$ and $\text{Out } F_n$ are called the inner automorphism group and the outer automorphism group of the free group respectively. We define a group IO_n to be the

quotient group of IA_n by $\text{Inn } F_n$. The group IO_n is the kernel of the natural map $\bar{\rho} : \text{Out } F_n \rightarrow GL(n, \mathbb{Z})$ induced by ρ . It is also known that the abelianization IO_n^{ab} of IO_n is given by $IO_n^{\text{ab}} \simeq (H^* \otimes_{\mathbb{Z}} \Lambda^2 H)/H$. (See [8, Theorem 6.2].) For $n \geq 2$ and $d \geq 2$, we define $IO_{n,d}$ to be the quotient group of $IA_{n,d}$ by $\text{Inn } F_n$. The group $IO_{n,d}$ is the kernel of $\pi_d \circ \bar{\rho}$. The second aim of this paper is to determine the structure of the abelianization $IO_{n,d}^{\text{ab}}$. The result is

Theorem 1.2. *For $n \geq 2$ and $d \geq 2$,*

$$IO_{n,d}^{\text{ab}} \simeq (IO_n^{\text{ab}} \otimes_{\mathbb{Z}} \mathbb{Z}/d\mathbb{Z}) \bigoplus \Gamma(n, d)^{\text{ab}}.$$

Finally, in Section 5, we compute the integral homology groups of $IA_{2,p}$ for an odd prime p . In general, to compute the integral homology groups of $IA_{n,d}$ is quite difficult as well as that of IA_n . In the case where $n = 2$ and $d = p$, we can compute those as follows:

Theorem 1.3. *For any prime p ,*

$$H_q(IA_{2,p}, \mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } q = 0, \\ \mathbb{Z}^{\oplus \alpha(p)} \oplus (\mathbb{Z}/p\mathbb{Z})^{\oplus 2} & \text{if } q = 1, \\ \mathbb{Z}^{\oplus (2\alpha(p)-2)} & \text{if } q = 2, \\ 0 & \text{if } q \geq 3 \end{cases}$$

where $\alpha(p) = 1 + \frac{(p-1)p(p+1)}{12}$ is the rank of $\Gamma(2, p)$ as a free group.

We remark that $IO_{2,p}$ is isomorphic to the congruence subgroup $\Gamma(2, p)$ since $IA_2 = \text{Inn } F_2$ due to Nielsen [12], and hence, it is a free group of rank $\alpha(p)$.

2. PRELIMINARIES

In this section we review the IA-automorphism group of a free group and the first Johnson homomorphism of the automorphism group of a free group. Throughout this paper we use the following notation and conventions.

- The group $\text{Aut } F_n$ acts on F_n from the right.
- For any $\sigma \in \text{Aut } F_n$ and $x \in F_n$, the action of σ on x is denoted by x^σ .
- For elements x and y of a group, the commutator bracket $[x, y]$ of x and y is defined to be $[x, y] := xyx^{-1}y^{-1}$.

2.1. The IA-automorphism group.

In this subsection, we prepare generators of IA_n , and some basic exact sequences which is required to prove our main theorems.

Let F_n be a free group on $\{x_1, \dots, x_n\}$. Magnus [11] showed that IA_n is finitely generated by automorphisms

$$K_{ij} : \begin{cases} x_i & \mapsto x_j^{-1} x_i x_j, \\ x_t & \mapsto x_t, \end{cases} \quad (t \neq i)$$

for distinct $i, j \in \{1, 2, \dots, n\}$ and

$$K_{klm} : \begin{cases} x_k & \mapsto x_k x_l x_m x_l^{-1} x_m^{-1}, \\ x_t & \mapsto x_t, \end{cases} \quad (t \neq k)$$

for distinct $k, l, m \in \{1, 2, \dots, n\}$ such that $l < m$. Since IO_n is the quotient group $IA_n/\text{Inn } F_n$, IO_n is also generated by (the coset classes of) automorphisms K_{ij} and K_{ijk} .

Next we give some basic exact sequences. Since the natural maps ρ and $\bar{\rho}$ are surjective, for any $n \geq 2$ and $d \geq 2$, we have exact sequences

$$(1) \quad 1 \rightarrow IA_n \rightarrow IA_{n,d} \xrightarrow{\rho} \Gamma(n, d) \rightarrow 1$$

and

$$(2) \quad 1 \rightarrow IO_n \rightarrow IO_{n,d} \rightarrow \Gamma(n, d) \rightarrow 1$$

respectively. Furthermore, by definition, we have

$$(3) \quad 1 \rightarrow \text{Inn } F_n \rightarrow IA_{n,d} \rightarrow IO_{n,d} \rightarrow 1.$$

These exact sequences are used in later sections.

2.2. The first Johnson homomorphism.

In this subsection, we review the first Johnson homomorphism of the automorphism group of a free group. For each $k \geq 1$, let $\Gamma_n(k)$ be the k -th subgroup of the lower central series of F_n defined by

$$\Gamma_n(1) := F_n, \quad \Gamma_n(k) := [\Gamma_n(k-1), F_n], \quad k \geq 1.$$

We denote the graded quotients $\Gamma_n(k)/\Gamma_n(k+1)$ by $\mathcal{L}_n(k)$. Set $\mathcal{L}_n = \bigoplus_{k \geq 1} \mathcal{L}_n(k)$. Then it is well known that \mathcal{L}_n naturally has a structure of a graded Lie algebra over \mathbb{Z} induced from the commutator bracket on F_n and it is naturally isomorphic to a graded free Lie algebra over H . In particular, we have $\mathcal{L}_n(1) = H$ and $\mathcal{L}_n(2) = \Lambda^2 H$. (For details, see [14].)

In this paper, for any $x \in \Gamma_n(k)$, we also denote by x the coset class of x in $\mathcal{L}_n(k)$.

For each $k \geq 1$, let

$$\tau' : IA_n \rightarrow \text{Hom}_{\mathbf{Z}}(H, \mathcal{L}_n(2)) = H^* \otimes_{\mathbf{Z}} \Lambda^2 H$$

be the homomorphism defined by

$$\sigma \mapsto (x \mapsto x^{-1}x^\sigma).$$

for $\sigma \in IA_n$ and $x \in H$. The map τ' naturally induces a homomorphism

$$\tau : IA_n^{\text{ab}} \rightarrow H^* \otimes_{\mathbf{Z}} \Lambda^2 H.$$

These homomorphisms τ' and τ are called the first Johnson homomorphisms of $\text{Aut } F_n$. The map τ is a $GL(n, \mathbf{Z})$ -equivariant isomorphism. In particular, IA_n^{ab} is a free abelian group of rank $\frac{1}{2}n^2(n-1)$. (See [8, Theorem 6.1].)

Next, we consider IO_n^{ab} . For any $y \in F_n$, we denote by ι_y the inner automorphism of F_n such that $x^{\iota_y} = y^{-1}xy$ for any $x \in F_n$. Considering a natural isomorphism $\text{Inn } F_n \rightarrow F_n; \iota \mapsto y$, we often identify $\text{Inn } F_n$ with F_n . Then we have $(\text{Inn } F_n)^{\text{ab}} = H$. It is also known that the induced homomorphism $(\text{Inn } F_n)^{\text{ab}} = H \rightarrow H^* \otimes_{\mathbf{Z}} \Lambda^2 H = IA_n^{\text{ab}}$ from the inclusion map $\text{Inn } F_n \hookrightarrow IA_n$ is injective, and whose image, which we identify with H by this map, is a direct summand of $H^* \otimes_{\mathbf{Z}} \Lambda^2 H$ as a \mathbf{Z} -module. Hence we see IO_n^{ab} is isomorphic to a free abelian group $(H^* \otimes_{\mathbf{Z}} \Lambda^2 H)/H$ of rank $\frac{1}{2}n(n+1)(n-2)$. For details, see [8, Theorem 6.2].

Now, the first Johnson homomorphism τ' induces a homomorphism

$$\tau'_d : IA_n \rightarrow (H^* \otimes_{\mathbf{Z}} \Lambda^2 H) \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z}.$$

Finally, we recall that τ'_d is extended to a homomorphism from $IA_{n,d}$ to $(H^* \otimes_{\mathbf{Z}} \Lambda^2 H) \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z}$. For any $\mathbf{Z}/d\mathbf{Z}$ -valued Magnus expansion θ , Kawazumi [8] constructed a crossed homomorphism

$$\tau^\theta : \text{Aut } F_n \rightarrow (H^* \otimes_{\mathbf{Z}} \Lambda^2 H) \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z}$$

and showed that if we restrict it to $IA_{n,d}$, then the map τ^θ is a homomorphism. Furthermore he also show that $\tau^\theta \equiv \tau'_d$ on IA_n . Especially the restriction $\tau^\theta|_{IA_n}$ is independent of the choice of the Magnus expansion θ . For details, see [8, Theorem 3.1].

3. THE ABELIANIZATION OF $IA_{n,d}$.

In this section we give a proof of Theorem 1.1. First, we see that since the first Johnson homomorphism τ is a $GL(n, \mathbf{Z})$ -equivariant isomorphism, τ induces a surjective homomorphism

$$\tilde{\tau} : H_0(\Gamma(n, d), IA_n^{\text{ab}}) \rightarrow (H^* \otimes_{\mathbf{Z}} \Lambda^2 H) \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z}.$$

To show $\tilde{\tau}$ is an isomorphism, we use

Lemma 3.1. *For $n \geq 2$ and $d \geq 2$, we have*

$$d[K_{ij}] = 0 \quad \text{and} \quad d[K_{klm}] = 0$$

in $H_0(\Gamma(n, d), IA_n^{\text{ab}})$.

Then considering the homological five term exact sequence

$$\begin{aligned} H_2(IA_n, \mathbf{Z}) \rightarrow H_2(IA_{n,d}, \mathbf{Z}) \rightarrow H_0(\Gamma(n, d), IA_n^{\text{ab}}) \\ \xrightarrow{\eta} IA_{n,d}^{\text{ab}} \rightarrow \Gamma(n, d)^{\text{ab}} \rightarrow 0. \end{aligned}$$

of (1), and a homomorphism

$$\tau^\theta : IA_{n,d} \rightarrow (H^* \otimes_{\mathbf{Z}} \Lambda^2 H) \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z}$$

induced from a Magnus expansion θ , we see η is injective. Hence we have a split exact sequence

$$0 \rightarrow (H^* \otimes_{\mathbf{Z}} \Lambda^2 H) \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z} \xrightarrow{\eta} IA_n^{\text{ab}} \rightarrow \Gamma(n, d)^{\text{ab}} \rightarrow 0.$$

This completes the proof of Theorem 1.1.

Here we consider the case where d equals to an odd prime integer p . For $n \geq 3$, Lee and Szczarba [10] showed that the abelianization $\Gamma(n, p)^{\text{ab}}$ of the congruence subgroup $\Gamma(n, p)$ is a $\mathbf{Z}/p\mathbf{Z}$ -vector space of dimension $n^2 - 1$. For $n = 2$, Frasch [5] showed that the congruence subgroup $\Gamma(2, p)$ is a free group of rank $\alpha(p) := 1 + \frac{(p-1)p(p+1)}{12}$. Furthermore Nielsen [12] showed that $IA_2 = \text{Inn } F_2$. Hence we have

Corollary 3.1.

$$IA_{n,p}^{\text{ab}} = \begin{cases} \mathbf{Z}^{\oplus \alpha(p)} \oplus (\mathbf{Z}/p\mathbf{Z})^{\oplus 2} & \text{if } n = 2, \\ (\mathbf{Z}/p\mathbf{Z})^{\oplus \frac{1}{2}(n-1)(n^2+2n+2)} & \text{if } n \geq 3. \end{cases}$$

4. THE ABELIANIZATION OF $IO_{n,d}$.

In this section we give a proof of Theorem 1.2. Considering the homological five term exact sequence of (3), we have

$$\begin{aligned} H_2(IA_{n,d}, \mathbf{Z}) \rightarrow H_2(IO_{n,d}, \mathbf{Z}) \rightarrow H_0(IO_{n,d}, (\text{Inn } F_n)^{\text{ab}}) \\ \xrightarrow{\delta} IA_{n,d}^{\text{ab}} \rightarrow IO_{n,d}^{\text{ab}} \rightarrow 0. \end{aligned}$$

Since H and $H^* \otimes_{\mathbf{Z}} \Lambda^2 H$ is free abelian groups, the injective homomorphism

$$H = (\text{Inn } F_n)^{\text{ab}} \hookrightarrow IA_n^{\text{ab}} = H^* \otimes_{\mathbf{Z}} \Lambda^2 H$$

induces an injective homomorphism

$$\psi_d : H \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z} \hookrightarrow (H^* \otimes_{\mathbf{Z}} \Lambda^2 H) \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z}.$$

For each i , $1 \leq i \leq n$, set $\iota_i := \iota_{x_i}$. Then, $\text{Inn } F_n$ is a free group on $\{\iota_1, \dots, \iota_n\}$.

Lemma 4.1. *For $n \geq 2$ and $d \geq 2$,*

$$d[\iota_i] = 0, \quad 1 \leq i \leq n$$

in $H_0(IO_{n,d}, (\text{Inn } F_n)^{\text{ab}})$.

Hence there exists an isomorphism

$$\bar{\xi} : H \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z} \rightarrow H_0(IO_{n,d}, (\text{Inn } F_n)^{\text{ab}})$$

such that $\psi_d = \delta \circ \bar{\xi}$, and we have a short exact sequence

$$0 \rightarrow H \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z} \xrightarrow{\delta} IA_{n,d}^{\text{ab}} \rightarrow IO_{n,d}^{\text{ab}} \rightarrow 0,$$

and hence

$$\begin{aligned} ((H^* \otimes_{\mathbf{Z}} \Lambda^2 H) \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z}) / (H \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z}) \\ \simeq ((H^* \otimes_{\mathbf{Z}} \Lambda^2 H)/H) \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z}, \\ \simeq IO_n^{\text{ab}} \otimes_{\mathbf{Z}} \mathbf{Z}/d\mathbf{Z}. \end{aligned}$$

This completes the proof of Theorem 1.2.

For $n \geq 2$ and an odd prime p , by an argument similar to that in Corollary 3.1, we obtain

Corollary 4.1.

$$IO_{n,p}^{\text{ab}} = \begin{cases} \mathbf{Z}^{\oplus \alpha(p)} & \text{if } n = 2, \\ (\mathbf{Z}/p\mathbf{Z})^{\oplus \frac{1}{2}(n+1)(n^2-2)} & \text{if } n \geq 3. \end{cases}$$

5. THE INTEGRAL HOMOLOGY GROUPS OF $IA_{2,p}$

In this section, we compute the integral homology groups of $IA_{2,p}$ for any odd prime p . Since the groups IA_2 and $\Gamma(2, p)$ are free groups stated above, considering the homological Lyndon-Hochschild-Serre spectral sequence of (1) for $n = 2$ and $d = p$, we see the homological dimension of $IA_{2,p}$ is 2. On the other hand, since the first homology group $H_1(IA_{2,p}, \mathbf{Z})$ is obtained in Section 3, it suffices to compute the second homology group $H_2(IA_{2,p}, \mathbf{Z})$. Our result is

Theorem 5.1. *For any odd prime p ,*

$$H_2(IA_{2,p}, \mathbf{Z}) = \mathbf{Z}^{\oplus (2\alpha(p)-2)}$$

where $\alpha(p) = 1 + \frac{(p-1)p(p+1)}{12}$.

To prove this theorem, first, we directly compute the second cohomology groups of $IA_{2,p}$. Then, using the universal coefficients theorem, we obtain the second homology group of $IA_{2,p}$.

Proposition 5.1. *For any odd prime integer p , we have*

$$H^2(IA_{2,p}, \mathbf{Z}) = \mathbf{Z}^{\oplus (2\alpha(p)-2)} \oplus (\mathbf{Z}/p\mathbf{Z})^{\oplus 2}.$$

Similarly, we obtain

Proposition 5.2. *For any odd prime integer p , we have*

$$H^2(IA_{2,p}, \mathbf{Z}/q\mathbf{Z}) \simeq \begin{cases} (\mathbf{Z}/q\mathbf{Z})^{\oplus (2\alpha(p)-2)} & \text{if } (q, p) = 1, \\ (\mathbf{Z}/q\mathbf{Z})^{\oplus (2\alpha(p)-2)} \oplus (\mathbf{Z}/p\mathbf{Z})^{\oplus 2} & \text{if } q = p^e. \end{cases}$$

Using Propositions 5.1 and 5.2, we obtain the second homology group $H_2(IA_{2,p}, \mathbf{Z})$ by the universal coefficients theorem.

6. ACKNOWLEDGMENTS

The author would like to express his sincere gratitude to Professor Nariya Kawazumi for his valuable advice and especially access to his unpublished work. He is also grateful to Professor Fred Cohen and Professor Benson Farb for explaining their unpublished works for the abelianization of the IA-automorphism group of a free group. Finally he would like to express his thanks to the referee for helpful comments and suggestions. The author is supported by JSPS Research Fellowships for Young Scientists.

REFERENCES

- [1] D. Arlettaz; On the homology and cohomology of congruence subgroups, J. of Pure and Applied Algebra 44 (1987), 3-12.
- [2] F. Cohen and J. Pakianathan; On Automorphism Groups of Free Groups, and Their Nilpotent Quotients, preprint.
- [3] F. Cohen and J. Pakianathan; On subgroups of the automorphism group of a free group and associated graded Lie algebras, preprint.
- [4] B. Farb; Automorphisms of F_n which act trivially on homology, in preparation.
- [5] H. Frasci; Die Erzeugenden der Hauptkongruenzgruppen für Primzahlstufen, Math. Ann. 108 (1933), 230-252.
- [6] S. M. Gersten; A presentation for the special automorphism group of a free group, J. of Pure and Applied Algebra 33 (1984), 269-279.
- [7] A. Hatcher and K. Vogtmann; Rational homology of $\text{Aut}(F_n)$, Math. Res. Lett. 5 (1998), 759-780.
- [8] N. Kawazumi; Cohomological aspects of Magnus expansions, preprint, The University of Tokyo. UTMS 2005-18 (2005), <http://xxx.yukawa.kyoto-u.ac.jp/abs/math.GT/0505497>.
- [9] S. Krstić and J. McCool; The non-finite presentability in $IA(F_3)$ and $GL_2(\mathbb{Z}[t, t^{-1}])$, Invent. Math. 129 (1997), 595-606.
- [10] R. Lee and R. H. Szczarba; On the homology and cohomology of congruence subgroups, Invent. Math. 33 (1976), 15-53.
- [11] W. Magnus; Über n -dimensionale Gittertransformationen, Acta Math. 64 (1935), 353-367.
- [12] J. Nielsen; Die Isomorphismen der allgemeinen unendlichen Gruppe mit zwei Erzeugenden, Math. Ann. 78 (1918), 385-397.
- [13] J. Nielsen; Die Isomorphismengruppe der freien Gruppen, Math Ann., 91 (1924), 169-209.
- [14] C. Reutenauer; Free Lie Algebras, London Mathematical Society monographs, new series, no. 7, (Oxford University Press, 1993).

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA
MEGURO-KU TOKYO 153-0041, JAPAN

E-mail address: takao@ms.u-tokyo.ac.jp